

CONTRIBUTION TO THE STUDY OF LITTORAL DIPTERANS I.  
ON A COLLECTION OF CERATOPOGONIDAE,  
CHIRONOMIDAE, PSYCHODIDAE AND  
TABANIDAE FROM SAGAR  
ISLAND, SUNDARBAN.

SANTANU RAY AND AMALESH CHOUDHURY\*

*Zoological Survey of India,  
Calcutta-700 053*

*\*Department of Marine Science  
University of Calcutta  
35 Ballygunge Circular Road  
Calcutta-700019.*

INTRODUCTION

This paper deals with the Dipteran insects collected during July, 1982 to December, 1983 from the intertidal zone of Ganga estuary, Sagar Island.

The families dealt with are Ceratopogonidae, Chironomidae, Psychodidae and Tabanidae.

Soils were collected at every fortnight from the intertidal zone and larvae and pupae were isolated in the laboratory of Susama Devichoudhurani Marine Biological Research Institute, Sagar Island by standardized floatation technique (Davies and Linley, 1966). After isolation, larvae and pupae were reared in the laboratory and adult Dipterans were collected.

SYSTEMATIC ACCOUNT

Order : DIPTERA

Suborder : NEMATOCERA

Family : CERATOPOGONIDAE

Tribe : **Ceratopogonini**

1. **Alluaudomyia maculosipennis** Tokunaga

1940. *Alluaudomyia maculosipennis* Tokunaga *Tenthredo* 3 : 181.

1942. *Alluaudomyia maculosipennis* Okada, *Trans. Nat. Hist. Soc. Formosa* 32 : 316 (Tiwan)

1959. *Alluaudomyia maculosipennis* Tok. and Mur. *Ins. Micronesia* 12 : 353 (Palau and Yap Is.)

1964. *Alluaudomyia maculosipennis* Wirth and Delfinado *Pacific Insect* 6 (4) : 627.

*Materials* : 4 ex (1 ♂ 3 ♀), Dhablat, Sagar South and Beguakhali, 19.ix.1984.

*Distribution* : Formosa, Java, Malaya, Philippines (Mindanao), Thailand, Viet Nam, Caroline Islands, Sri Lanka (Ceylon), India.

*Remark* : The species is being reported from India for the first time.

## 2. *Alluaudomyia formosana* Okada

1942. *Alluaudomyia maculipennis* (Carter, Ingran and Macfie) var. *formosana* Okada, Trans. Nat. Hist. Soc. Formosa 32 : 317.

*Materials* : 5 ex (5 ♀ ♀), Dhablat, Sagar South and Beguakhali, 19.ix.1984.

*Distribution* : Formosa, Malaya, Thailand, Sri Lanka (Ceylon) and India.

*Remark* : The species is being reported from India for the first time.

## Tribe : Culicoidini

### 3. *Culicoides oxystoma* Kieffer

1910. *Culicoides oxystoma* Kieffer, Mem. Indian. Mus. 2 : 193 (♀).

1913. *Culicoides kiefferi* Patton, Ind. J. Med. Res. 1 : 336 (♂ ♀ Pl ; Proce. Goetghebuer 1910).

1921. *Culicoides pattoni* Kieffer, Bull. Soc. Ent. Fr.

1921. (n name for kiefferi Patton)

1938. *Culicoides housei* Causey, Am. J. Hyg. 27 : 407 (♂).

1951. *Culicoides punctigerus* Tokunaga, Scient. Rep. Saikyo. Univ. (Agric) 1 : 101 (♂ ♀).

1956. *Culicoides alatus* Dasgupta and Ghosh, Bull, Cal. Sch. Trop. Med. Hyg. 4 : 162 (♀).

*Materials* : 7 ex (4 ♂ ♂, 3 ♀ ♀), Sagar South, Dhablat and Beguakhali, 19.ix.1984.

*Distribution* : India, Bangkok, Thailand, Batavia (Djakarta), Java.

### 4. *Culicoides peliliouensis* Tokunaga

1936. *Culicoides peliliouensis* Tokunaga, Mushi 9 : 55 (♀).

1961. *Culicoides ejercitoi* Delfinado, Fieldiana. zool. 33 : 643 (♀).

*Materials* : 21 ex (14 ♂ ♂, 7 ♀ ♀) Sagar South, Beguakhali 19.ix.84.

*Distribution* : Akarokuru, Peliliou Island, Palan, Sink, Calamianes (Cuilon) Island, Palawan, Philippines, India.

*Remark* : This species is also being reported from India for the first time.

5. **Culicoides similis** Carter, Ingran and Macfie.

1920. *Culicoides similis* Carter, Ingran and Macfie Ann. Trop. Med. Parasitol. 14 : 255 (♂ ♀).

**Materials** : 4 ex (2 ♂ ♂, 2 ♀ ♀), Dhablat, Sagar South and Beguakhali, 19.ix.1984.

**Distribution** : India, Laos, Malaya, Thailand and Africa.

Tribe : **Sphaeromiini**6. **Nilobezzia acanthopus** (de Meijere)

1907. *Ceratopogon* (*Bezzia*) *acanthopus* de Meijere Tijdschr. Ent. 50 : 215 (♀).

1931. *Bezzia* (*Nilobezzia*) *acanthopus* (de Meijere) Johannsen 439.

1931. *Bezzia* (*Nilobezzia*) *raphaelis* var. *conspicua* Johannsen 411.

1934b. *Nilobezzia conspicua* (Johannsen) 441.

**Materials** : 4 ex (4 ♀ ♀) Dhablat, Sagar South and Beguakhali, 19.ix.1984.

**Distribution** : Java, Sumatra, Thailand, India.

**Remarks** : The species is being reported from India for the first time.

Family : **CHIRONOMIDAE**

Tribe : **Chironomini**

7. **Chironomus barbatitarsis** Kieffer

1911. *Chironomus barbatitarsis* Kieffer, Rec. Indian. Mus 6 : 154.

**Materials** : 4 ex (2 ♂ 2 ♀) Dhablat, Beguakhali and Sagar South.

**Distribution** : Burma, India.

Superfamily : **PSYCHODOIDEA**

Family : **PSYCHODIDAE**

Subfamily : **PSYCHODINAE**

8. **Psychoda nigripennis** Brunetti

1908. *Psychoda nigripennis* Brunetti, Rec. Indian. Mus. 2 : 376 (♂ ♀).

**Material** : 4 ex (2 ♂ 2 ♀) Dhablat, Beguakhali and Sagar South, 6.iii.1983.

**Distribution** : India.

Suborder : BRACHYCERA

Family : TABANIDAE

Subfamily : TABANINAE

Tribe : **Tabanini**

### 9. *Tabanus striatus* Fabricius

1778. *Tabanus striatus* Fabricius, Mantissa Insect 2 : 256 (sex ?).

1827?. *chinensis* Thunberg Nova. Acta. Soc. Sci. Upsal 9 : 61 (? sex).

1842?. *dorsilinea* Wiedemann, Analecta Ent. ?? (♂).

1850. *Tabanus hilaris* Walker, Insecta Saundersiana 1 : 49 (♂).

*Materials* : 4 ex, (3♂ 1♀), Dhablat, Beguakhali and Sagar South (Sex ?)

*Distribution* : Ceylon, China, India, N. Thailand, N. Viet Nam. W. Pakistan and Africa.

### 10. *Atylotus agrestis* (Wiedemann)

1828. *Atylotus agrestis* (Wiedemann) Aussereurop. zweifl. Insekt. 1 : 557 (♂ *Tabanus*).

1838. *Atylotus ditoeniatus* Maequart, Dept. exot. 1 (1) : 126 (130) (♂ ♀ as *Tabanus ditoeniatus*, p. 215 (219)).

1850. *Atylotus pyrrhus* Walker, Insecta Saundersiana 1 : 47 (♂ ♀ *Tabanus*).

1868. *Atylotus agricola* of Schiner, not Wiedemann, mis identification.

### SUMMARY

The present paper is based on a collection of Dipterans from littoral zone of the Ganga estuary. Six species of Ceratopogonidae, one species of Chironomidae, one species of Psychodidae and two species of Tabanidae are reported in this paper. Four species viz. *Alluaudomyia maculosipennis* Tokunaga, *Alluaudomyia formosana* Okada *Culicoides peliliouensis* Tokunaga and *Nilobezzia acanthopus* (de Meijere) are being recorded from India for the first time.

### ACKNOWLEDGEMENT

One of the authors Santanu Ray is grateful to the Director, Zoological Survey of India, Calcutta, for the necessary permission to collect the material from Sagar Island, Sundarban. We are thankful to Dr. W. W. Wirth, Dr. M. Dutta, Dr. P. Choudhuri and Mr. P. Parui for their kind help in the identification of Ceratopogonidae, Tabanidae, Chironomidae and Psychodidae respectively. We are also thankful to the authorities of Susama Devichoudhurani Marine Biological Research Institute for the facilities provided.

## REFERENCES

- DAVIES, J. B. AND LINLEY, J. R. (1966). A standardized floatation method for separating *Leptoeonops* (Diptera : Ceratopogonidae) and other larvae from sand samples. *Mosquito News* 26 : 440.
- DELFINADO, M. D. AND D. E. HARDY. 1973. *A catalog of the Diptera of the Oriental Region. Vol. I Suborder Nematocera. The University Press of Hawaii, Honolulu.*
- DELFINADO, M. D. AND D. E. HARDY, 1973. *A catalog of the Diptera of the Oriental Region. Vol. II The University Press of Hawaii, Honolulu.*
- WIRTH, W. W. AND MERCEDES D. DELFINADO. 1964 Revision of the oriental species of *Alluaudomyia* Kieffer (Diptera, Ceratopogonidae), *Pacific Insects* 6 (4) : 599-648.
- WIRTH, W. W. AND NIPHAN C. RATANAWARABHAN, 1981. New species and records of predaceous midges (Diptera : Ceratopogonidae) from Rice Paddies in Thailand. *Pacific Insects* 23 (3-4-) 396-431.