

IX. CONTRIBUTIONS TO THE FAUNA OF  
YUNNAN BASED ON COLLECTIONS  
MADE BY J COGGIN BROWN,  
B Sc , 1909—1910 <sup>1</sup>

PART VII.—TABANIDAE.

By GERTRUDE RICARDO.

The Tabanidae sent me for identification by Dr. Annandale are few in number, comprising the following species:—

**TABANUS**, Linné

**Tabanus orientis**, Walker. Two females from Man Hsien.

**Tabanus monotaeniatus**, Bigot. Two females from the same locality, rather smaller in size than usual, the abdomen being narrower, and reddish in colour.

**HAEMATOPOTA**, Meigen.

**Haematopota lata**, Ricardo. Four females from Man Hsien.

**CHRYSOPS**, Meigen.

**Chrysops mlokosiewiczi**, Bigot (*Chrysops striatus*, Wulp). A series of females from between Tengyueh and Tali Fu (J Coggin Brown). The frontal callus is reddish yellow, not black. This species has a wide distribution in the Palaearctic and Oriental Regions, being recorded from China, Japan and Persia.

**Chrysops designata**, Ricardo (p. 383, *antea*). One female.

For references to these species see *Rec. Ind. Mus.*, vol. iv, No. vi, 1911, containing a revision of the species of *Tabanus* from the Oriental Region, and the same publication, vol. iv, No. viii, for species of *Chrysops* and other genera of Tabanidae.

---

<sup>1</sup> The previous papers of this series appeared in *Rec. Ind. Mus.*, v, p. 193 (1910), and vi, pp. 13 and 215 (1911).