

PART V.—BIBIONIDAE, MYCETOPHILIDAE AND ANOPHELINAE.

[Among the Diptera brought back by Mr. Coggin Brown there are few Nematocera. Two species of Bibionidae and one of Mycetophilidae have been identified by Mr. Brunetti and two of Anopheline mosquitoes by Major S. P. James, I.M.S., as follows:—]

BIBIONIDAE.

Plecia melanaspis, Wied.

Localities.—Mong-Wan; Lahsa; between Tengyueh and Tali-Fu.

[A common Himalayan species the range of which extends to eastern China and the Malay Archipelago.—N. A.]

Dilophus graciosus, Big.

Locality.—Between Tengyueh and Tali-Fu.

[Not uncommon in the Himalayas, along the whole of which it is apparently found.—N. A.]

MYCETOPHILIDAE.

Sciara rufithorax, v. Wulp.

Locality.—Tengyueh.

[Common almost all over India and in many parts of Malaysia.—N. A.]

ANOPHELINAE.

Nyssorhynchus maculipalpis, James and Liston.

Localities.—Chu-Chih and Lahsa.

[According to Theobald (*Mon. Culic.*, v, p. 62, 1910) this is *N. indiensis*, Theob., which he now regards as a distinct species. It occurs in W. and Central India. Theobald's name has priority.—N. A.]

Myzorhynchus sinensis, Wied.

Localities.—Chu-Chih and Man-Nau.

[This species is apparently common in India, but its synonymy is very doubtful.—N. A.]