



VARIATION OF MEIOFAUNA DURING DAY TIME FOLLOWING FULL MOON AND NEW MOON

G. SIVALEELA AND K. VENKATARAMAN*

Marine Biology Regional Centre, Zoological Survey of India
130, Santhome High Road, Chennai-600 028

*Zoological Survey of India, M-Block, New Alipore, Kolkata-700 053
venkyzsi56@yahoo.com

ABSTRACT

Aim of this research was to investigate meiofaunal movement patterns in day time at two stations in the polluted and unpolluted area in every three hours during full moon and new moon time during 2006 & 2007. Meiofauna was abundant in all samples, the animals are Harpacticoida, Nematoda, Oligochaeta, Tardigrada, Ostracoda etc.

INTRODUCTION

Meiofauna variation during sunlight period was studied at two stations namely station 1

(Adayar) stressed by sewage pollution and station 2 (Besant Nagar) a relatively unpolluted area (Tables 4 a,b). Meiofauna population in the two stations was composed of nematodes, harpacticoid copepods, foraminiferans, polychaetes, oligochaetes and ostracods.

The nematodes comprised 25-40% of the population at station 1 and 53-65% at station 2 (Fig. 1 a, b). Nematodes, copepods and foraminiferans formed more than 60% at station 1 and 80% at station 2, and small percentage was made up by communities of polychaetes and ostracods (Fig. 1 a, b).



Table 1a. Variation of meiofauna density at Adayar following New Moon (Station 1) during 2006 and 2007.

Fauna	Time intervals									
	6am		9am		12pm		3pm		6pm	
	2006	2007	2006	2007	2006	2007	2006	2007	2006	2007
Nematodes	160	112	148	120	210	160	132	120	120	80
Copepods	40	34	35	48	42	25	30	35	25	12
Foraminiferans	15	40	87	59	75	30	60	44	35	18
Polychaetes	65	38	98	25	75	45	53	43	25	30
Oligochaetes	120	90	180	129	194	140	100	110	80	102
Ostracodes	5	10	38	12	93	14	42	18	15	10
Total	405	324	586	393	689	414	417	370	300	252

Table 1b. Variation of meiofauna density at Besant Nagar following Full Moon (Station 2) during 2006 and 2007.

Fauna	Time intervals									
	6am		9am		12pm		3pm		6pm	
	2006	2007	2006	2007	2006	2007	2006	2007	2006	2007
Nematodes	220	350	210	378	350	410	310	390	250	250
Copepods	120	180	110	202	190	240	190	210	120	125
Foraminiferans	87	109	70	147	98	190	90	120	80	80
Polychaetes	25	50	20	50	25	98	30	44	20	40
Oligochaetes	15	20	10	12	10	15	12	22	10	15
Ostracodes	15	10	20	45	15	60	10	40	10	25
Total	482	719	440	834	688	1013	642	826	490	535



The mean density of meiofauna at station 1 was in the range of 300-689 n/10cm² and 252-414 n/10cm² during 2006 and 2007 respectively. At station 2, it was observed in the range of 440-688 n/10cm² and 535-1013 n/10cm² during 2006 and 2007 respectively. The nematode/Copepod-ratio (N/C-ratio) varied between 6:1 (station 1) and 2:1 (station 2). The wide range of N/C ratio was observed at station 1 (Adayar), probably due to pollution.

Fig. 1a. Mean percentage composition of meiofauna communities at Adayar and Besant Nagar 2006.**Fig. 1b.** Mean percentage composition of meiofauna communities at Adayar and Besant Nagar 2007.

With respect to variation during sunlight period, meiofauna was at its peak during midday (12 noon), probably due to availability of food (Fig. 2). Midday peaks in meiofauna generally indicate consumption of microalgae, but the diel response varies among taxa and demographic groups. Co-variation of environmental factors may complicate interpretation of feeding patterns during sunlight periods.

The abundance of meiofaunal groups is known to be controlled by physical factors like temperature, grain size, salinity, wave action, beach slope, tidal action and capillary water and its action as well.

Fig. 2. Mean total density of meiofauna at Adayar and Besant Nagar during 2006 and 2007.



ACKNOWLEDGEMENTS

The author is highly thankful to Director, ZSI, Kolkata for support and facilities provided to carryout my Ph.D programme and for his excellent guidance and encouragement in Ph.D and also thankful to Dr.C.Venkatraman Scientist-C and Officer-in-charge, MBRC/ZSI for his support.

REFERENCES

- Heininger, P., S. Hoss, E. Claus, J. Pelzer and Traunspurger, W. (2007). Nematode communities in contaminated river sediments. *Environ. Pollut.*, **146** : 64-76
- Pearson, T.H. and Rosenberg, R. (1978). Macrobenthic succession in relation to organic enrichment and pollution of the marine environment. *Oceanog. Mar. Biol. Ann. Rev.*, **16** : 229-311;
- Platt, H.M., K.M. Shaw and Lamshead, P.J.D. 1996. Nematode species abundance patterns and their use in the detection of environmental perturbations. *Hydrobiologia*, **118** : 59-66 (1984).
- Otway, N.M., C.A. Gray, J.R. Craig, T.A. Mcvea and Ling, J.E. Assessing the impacts of deepwater sewage outfalls on spatially-and temporally variable marine communities. *Mar. Environ. Res.*, **41** : 45-71.
- Shiddamallayya, N. and Pratima, M. (2008). Impact of domestic sewage on fresh water body. *J. Environ. Biol.*, **29** : 303-308
- Khurana, M.P.S. and Bansal, R.L. (2008). Impact of sewage irrigation on speciation of nickel in soils and its accumulation in crops of industrial towns of Punjab. *J. Environ. Biol.*, **29** : 793-798.